

## Thermodynamic Approach of Lanthanide-Iron Double Oxides Formation

**Authors :** Vera Varazashvili, Murman Tsarakhov, Tamar Mirianashvili, Teimuraz Pavlenishvili, Tengiz Machaladze, Mzia Khundadze

**Abstract :** Standard Gibbs energy of formation  $\Delta G_{\text{for}}(298.15)$  of lanthanide-iron double oxides of garnet-type crystal structure  $R_3\text{Fe}_5\text{O}_{12}$  - RIG (R - are rare earth ions) from initial oxides are evaluated. The calculation is based on the data of standard entropies  $S_{298.15}$  and standard enthalpies  $\Delta H_{298.15}$  of formation of compounds which are involved in the process of garnets synthesis. Gibbs energy of formation is presented as temperature function  $\Delta G_{\text{for}}(T)$  for the range 300-1600K. The necessary starting thermodynamic data were obtained from calorimetric study of heat capacity - temperature functions and by using the semi-empirical method for calculation of  $\Delta H_{298.15}$  of formation. Thermodynamic functions for standard temperature - enthalpy, entropy and Gibbs energy - are recommended as reference data for technological evaluations. Through the isostructural series of rare earth-iron garnets the correlation between thermodynamic properties and characteristics of lanthanide ions are elucidated.

**Keywords :** calorimetry, entropy, enthalpy, heat capacity, gibbs energy of formation, rare earth iron garnets

**Conference Title :** ICCEAC 2015 : International Conference on Chemical Engineering and Applied Chemistry

**Conference Location :** Rome, Italy

**Conference Dates :** September 17-18, 2015